

ODD PROTOCOL

---

## **ODD protocol of the FarmECS-IBR model**

A model to support decision-making in the context of a national  
IBR eradication program

---

This model was developed by  
ECOEPI

Department of Ecological Modelling  
Helmholtz Centre for Environmental Research - UFZ

*1st Developer*  
Dr. Martin Lange

*2nd Developer*  
Jonas Brock

*3rd Developer*  
Dr. Hans-Hermann Thulke

December 13, 2020

# Table of Contents

---

Table of Contents	I
List of Figures	II
List of Tables	II
List of Acronyms	II
<b>1 ODD protocol of the FarmECS-IBR model</b>	<b>1</b>
1.1 Overview . . . . .	1
1.1.1 Purpose . . . . .	1
1.1.2 Entities, state variables and scales . . . . .	1
1.1.3 Process overview and scheduling . . . . .	3
1.2 Design . . . . .	3
1.2.1 Herd management processes . . . . .	3
1.2.2 Animal transports . . . . .	5
1.2.3 Epidemiological processes . . . . .	6
1.3 Details . . . . .	6
1.3.1 Globals . . . . .	6
1.3.2 Input data . . . . .	7
1.3.3 Initialisation . . . . .	7
1.3.4 Submodels . . . . .	8
1.3.4.1 Replacement . . . . .	8
1.3.4.2 Grouping . . . . .	9
1.3.4.3 Breeding . . . . .	9
1.3.4.4 Abortion . . . . .	10
1.3.4.5 Calving . . . . .	11
1.3.4.6 Transports . . . . .	11
1.3.4.7 Infection . . . . .	12
1.3.4.8 Mortality . . . . .	15
1.3.4.9 Virus release . . . . .	15
<b>Bibliography</b>	<b>16</b>

## List of Figures

---

Figure 1.1:	Structure of the FarmECS-IBR model . . . . .	2
Figure 1.2:	Typical animal transports under the Irish cattle management system . . .	6
Figure 1.3:	Epidemiological states in the model . . . . .	13

## List of Tables

---

Table 1.1:	Globally accessible parameters . . . . .	7
Table 1.2:	Data used to describe the Irish cattle system . . . . .	7
Table 1.3:	Parameters relevant for model initialisation . . . . .	8
Table 1.4:	Model parameters related to replacement . . . . .	9
Table 1.5:	Model parameters related to abortion . . . . .	11
Table 1.6:	Model parameters related to calving . . . . .	11
Table 1.7:	Model parameters related to mortality . . . . .	15

## List of Acronyms

---

<b>BoHV-1</b>	Bovine Herpes Virus Type 1
<b>IBR</b>	Infectious Bovine Rhinotracheitis
<b>EU</b>	European Union
<b>ODD</b>	Overview, Design Concepts, and Details
<b>AIM</b>	Animal Identification and Movement
<b>DAFM</b>	Department of Agriculture, Food and the Marine
<b>SOM</b>	Self Organizing Maps

# 1 | ODD protocol of the FarmECS-IBR model

---

## 1.1 Overview

A stochastic large scale individual-based model of Bovine Herpes Virus Type 1 (BoHV-1) was developed in the framework of this contribution. The model, called FarmECS-Infectious Bovine Rhinotracheitis (IBR) is build up on the system understanding of the Irish cattle sector. This protocol provides a complete technical documentation of the developed model. The documentation of the model follows the Overview, Design Concepts, and Details (ODD) protocol a standardized scheme designed to produce a transparent and comprehensive model description following a generic structure ([Grimm et al., 2006, 2010, 2020](#)).

The FarmECS-IBR model is a compilation of a spatially explicit, stochastic, individual-based cattle herd-management and animal-trade model for the entire cattle sector of a geographic area. Superimposed to the herd-management and animal-trade model is a transmission and disease course model of BoHV-1 that represents the spread of IBR within and between herds.

### 1.1.1 Purpose

Purpose of the FarmECS-IBR model is the evaluation of different BoHV-1 control strategies to provide policy support and to assist with decision-making in the context of an Irish BoHV-1 eradication program.

### 1.1.2 Entities, state variables and scales

Spatial scale of the model is the Irish county Kerry located in the south-east of Ireland, with an area of approx.  $5'000 \text{ km}^2$ , 5'000 cattle herds and 360'000 individual cattle. The model runs in weekly timesteps and is individual-based, meaning that each animal is simulated explicitly.

The model consists of two entities, namely the animal and the herd entity (see Figure 1.1). Each entity is described by one or a set of state variables. Due to the application of an Entity-Component-System (ECS), state variables are grouped into Components that are attached to the entities. Components, and thus state variables, can be added to an entity or removed, depending on whether they are currently used or not.




Figure 1.1: Entities, components and state variables of the FarmECS-IBR model.

In the model, individual animals are the most complex entities. Each individual animal comprises at least an *AnimalBase* component that provides information on typical demographic variables such as an animal's age, sex, breed, date of death or membership in a management cohort. In addition each animals holds an *AnimalGroup* component specifying an animal's herd. The other components are optional and are attached to certain animals only temporarily. For example, female animals that have reached a specific age or are intended for breeding get the *AwaitsBreeding* component attached. This component stores all state variables necessary for breeding, such as the number of insemination trials, the last insemination trial, whether an insemination was successful (*inseminationSuccess*) and, if so, the date of conception (*dateConception*). If breeding was successful, the *AwaitsBreeding* component is detached and the *IsPregnant* component is attached. Among other things, this component provides information about the sex or race of the fetus. The components *ScheduledForSale*, *IsForSale* and *AnimalTradeSource* are used to handle animal transports between the simulated herds. The *AnimalBohvEpiStat* component keeps information on the animal's BoHV-1 infection status and when it last changed.

The herd components listed in Figure 1.1 are permanent for all cattle herds in the model. The

*HerdBase* component provides information on the ID of the herd, the herd type and lists all present animals. The *Position* component gives information on the spatial location of each herd and is used to determine neighbouring herds which is important for the epidemiological part of the model. Important herd management variables and parameters are stored in the *HerdManagement* component. This component describes how the farmer manages his herd (e.g. when he is breeding his animals or how many heifers he keeps annually to replace older cows). The *HerdBohvStatus* component stores herd BoHV-1 information, such as the number of animals in respective BoHV-1 health states or the total number of seropositive animals.

### 1.1.3 Process overview and scheduling

The FarmECS-IBR model runs in discrete timesteps of one week. At each timestep the main processes of the model are executed in the following order and animal- and herd-level state variables are updated wherever necessary:

- 1) Herd management: grouping of animals to cohorts, marking animals for sale
- 2) Breeding
- 3) Abortion
- 4) Calving
- 5) Trade/transport
- 6) BoHV-1 infection, recovery, reactivation
- 7) Background mortality/culling
- 8) Virus release

## 1.2 Design

According to [Brock et al. \(2020\)](#) the structure of the characteristics of cattle disease models can be described using three main features: [1] biological processes, [2] farming-related processes and [3] pathogen-related processes. In the FarmECS-IBR model, biological processes such as ageing, pregnancy, lactation, fertility and mortality are simulated at each time step for all individual animals. At herd level, management processes such as grouping, breeding, buying and selling of animals are simulated for each herd type. Independent of the biological- and management-related processes, pathogen-related processes are represented in the model. Thereby, each animal is assigned an epidemiological status, which can change according to individual infection or recovery.

### 1.2.1 Herd management processes

A detailed understanding of herd types and herd management is needed for animal disease control and surveillance activities, to inform epidemiological study design and interpretation,

and to guide effective policy decision-making. Therefore, sixteen different herd types are defined in the FarmECS-IBR model, representing the typical management schemes of the cattle sector in Kerry/Ireland. The herd types were identified from a previous analysis where expert knowledge was coupled with a machine-learning algorithm called Self Organizing Maps (SOM) to explore multi-dimensional herd data extracted from the Irish Animal Identification and Movement (AIM) database.

For the sake of model development, we have assigned each of the 5000 herds in Kerry to one of our identified herd types based on a classification scheme we developed for the Irish cattle population. In the model we have implemented a rule-based management system for each of the identified herd types, which is followed by all herds of the respective type. This management scheme determines, for example, how long herds of a certain type keep youngstock, at what age they sell them and whether herds maintain themselves through breeding. For each herd, its population structure emerges from a combination of biological parameters of individual animals and different management practices of herds. To ensure validity of the model, each herd types population structure is systematically compared to data extracted from the AIM livestock register.

In the following, a brief description of the identified/implemented herd types is given. Among the dairy herds, we identified four sub-groups that differ in terms of their management practices. Typical dairy herds (**D**) that sell their male calves (to **Sdm** herds) at the age of a few weeks are the most prominent dairy herd type in Kerry. In these herds, female calves are kept and reared as replacements. Non-rearing dairy herds (**DnR-C**) sell most of their calves, but female calves are moved to external contract rearing herds (**Rdf**). Most of them return to the **DnR-C** herd as pregnant heifers before calving. A similar herd type is the **DnR-nC** herd type. There are similarities to the previous herd type, with the sale of all new-born calves within a few weeks after birth and the introduction of replacement heifers. In contrast to the **DnR-C** herds, however, replacement heifers are not purchased from contract rearing herds, but from other herds with a surplus of such animals. There are also dairy herds (**DRm**) that keep all their calves, including their males calves until they are sold as youngstock for fattening to fattening (**F**) herds.

Five different beef production systems were identified from the data and implemented in the model. The first type we extracted is the beef pedigree herd (**BP**) whose main production objective is breeding of purebred beef cattle. Purebred beef herds are important for providing breeding stock (e.g. Beef bulls) to other commercial cattle producers in both the dairy and beef sectors. The suckling to beef (**BSB**) system includes the full beef production cycle, from birth through to the age of slaughter. Typically, calves stay with their dams until weaning at 6-8 months. Then, the weaned calves are retained for rearing and fattening in the same enterprise. The beef suckling to youngstock (**BSY**) system is very similar to the **BSB** system, however these herds do not fatten their cattle intended for slaughter. Instead these animals are mainly sold to fattening herds as yearlings (youngstock for fattening). Non-rearing suckling to youngstock (**BSY-nR**) herds are a variation of the **BSY** herd type, with the difference being that all female calves are sold after weaning. These herds mainly purchase pregnant heifers for replacement from other suckling herds or **Sbf** herds. Beef suckler to weaning (**BSW**) herds are

the most common beef system in Ireland. These herds sell most of their calves after weaning between six and eight months of age to store or fattening herds. A proportion of females are kept for replacement.

The store/rearing group mainly comprises herds that are non-calving. Overall, five sub-types (**Sdm**, **Rdf**, **Sbf**, **Sbm**, **Sbmx**) were identified and implemented, which differ in terms of their management practices. Store dairy bull (**Sdm**) herds typically purchase male calves between 10 and 30 days at age from dairy herds. These animals are reared and then sold to fattening herds. Three different types of beef store herds were identified, which differ in terms of their sex composition, rather than by management practices: store beef males (**Sbm**), store beef females (**Sbf**) and store beef mixed (**Sbmx**) systems. These herds purchase beef animals as weanlings and rear them until they are sent to fattening herds. In rearing dairy female (**Rdf**) herds, young female dairy calves are introduced, reared and inseminated, before being returned to their birth herd as pregnant heifers.

Herd which on average consist half pure-bred dairy animals and half animals cross-bred between dairy and beef were also identified and implemented. These herds produce milk on the one hand, but on the other hand they have another cattle enterprise, solely focused on beef production. We classified these dual purpose herds as mixed herds (**M**). Fattening herds (**F**) buy calves, weanlings and youngstock from a wide range of herd systems and fatten them until slaughter.

### 1.2.2 Animal transports

In the model animal movements emerge from supply and demand. Supply of and demand for animals by individual herds emerge from herd type specific management and stochastic processes, e.g. mortality or infertility.

Animals in supply and demand are classified into 7 trading cohorts, like e.g. youngstock for fattening or pregnant heifer dairy (see Figure 1.2). As an example, store dairy male herds (**Sdm**) herds have demand for male dairy calves (CDM) according to the free space in their premises. Dairy (D) herds supply male dairy calves (CDM) for rearing by trying to sell their male calves at an age of 2 weeks.

Animals intended to leave a herd are preferably distributed to enterprise sites and trade partners. If this is not possible, supplied animals enter an open market, where demanders select supplies based on distance as well as demand and supply batch size. If demand can't be satisfied on the market for a certain time, it is satisfied by imports from outside the modelled region.




Figure 1.2: Typical animal transports under the Irish cattle management system. For a better overview only primary movement cohorts are shown.

### 1.2.3 Epidemiological processes

Epidemiological processes of BoHV-1 infection are driven by direct transmission between animals in the same herd, by indirect transmission between nearby herds, as well as animal transports. On farm, transmission of the infection is independently modelled within management groups compared to transmission between groups.

Individual probability to contract infection follows standard epidemiological formalisms as described by Brock et al. (2020), comprising all pathways of the animal’s current exposure.

Reactivation of latently infected animals is explicitly modelled by means of a weekly probability. Infectiousness of primary infected animals is greater than of secondary infected animals.

## 1.3 Details

### 1.3.1 Globals

This section provides an overview on globally accessible data structures that are utilized by submodels to access global parameters. In the FarmECS-IBR model, globally available parameters are structured into two main groups: [1] biological parameters and [2] management-related parameters. Table 1.1 provides an overview of these parameters.

Table 1.1: Globally accessible parameters

	Parameter	Symbol	Type	Description	Value
Biological data	gestationLength	$\tau_{gest}$	int	Length of gestation (weeks)	XX
	lactationLength	$\tau_{lact}$	int	Length of lactation after calving (weeks)	XX
Management data	maxAgeNewborn	$a_{newb}$	int	Age to move from cohort newborn to calf	XX
	maxAgeCalf	$a_{calf}$	int	Age to move from cohort calf to heifer or grasser	XX
	maxAgeGrasser	$a_{gras}$	int	Age to move from cohort grasser to fattener	XX
	maxAgeFattener	$a_{fatt}$	int	Maximum age for animals in cohort fattener	XX
	minAgeBreeding	$a_{breed}$	int	Minimum age of heifers for first breeding	XX

### 1.3.2 Input data

The model does not require input data but in advance several datasets were analysed to provide an understanding of the Irish cattle system. Table 1.2 lists data used to describe and model the Irish cattle system. This table reflects the minimum information that would be needed to apply the model to other cattle populations.

Table 1.2: Data used to describe the Irish cattle system

Dataset	Description	Source
Herd population in 2017	List of all cattle registered in Ireland on 3 different dates (1st Jan, 1st May and 1st Sep) in 2017	AIM database
All births from 2016 to 2018	List of all births from 2016 to 2018	AIM database
All animal transports in 2017	List of all animal movements that took place in 2017	AIM database

Data listed in Table 1.2 was obtained from the AIM database maintained by the Department of Agriculture, Food and the Marine (DAFM). In accordance with European Union (EU) requirements, the AIM database comprises records on all births, movements and disposals (i.e. origin, destination and date), tracking each individual bovine in Ireland from birth or import to death or export (DAFM, 2017).

The herd population dataset, as well as the animal transport dataset were used to classify cattle herd types and to gain a detailed understanding of their management regimes. These datasets were also used to validate the model. The birth dataset was important to extract parameters relevant for breeding and calving.

### 1.3.3 Initialisation

The model is initialised by the following steps. First, all herds to be modelled are read from a table (herdTableFile, see Table 1.3) listing for each of the 5000 cattle herds in Kerry their position, herd type, target size, breeding and outdoor season. Then, for each herd in the table, animals are created and added to the engine. Important animal state variables, such as animals age, sex etc. are drawn from predetermined distributions. Afterwards, a 20-year pre-simulation

is conducted to ensure a proper initial population structure and valid animal components and state variable values.

After the 20-year pre-simulation, an individual herd's population is written to the herdXmlFile (see Table 1.3). This file lists all animals from all herds, along with its initialised state variables. At model start up, individual herd populations are initialized from the herdXmlFile by randomly drawing animals until the desired number of animals per herd (target size listed in herdTableFile) is reached.

Table 1.3: Parameters relevant for model initialisation

Parameter	Symbol	Type	Description
herdXmlFile	-	String	Path to xml file with serialized herd populations
herdTableFile	-	String	Path to herd table
columns...	-	String	Column names of the herd table file (e.g. ID, Type, Size)
poolsIndoor	$P_i(ij)$	List[Matrix[boolean]]	Per herd type pooling/contact matrix indoor
poolsOutdoor	$P_o(ij)$	List[Matrix[boolean]]	Per herd type pooling/contact matrix outdoor
maxRadius	$r_{neigh}$	float	Neighborhood radius

During initialization, a contact structure between the animals is defined for each herd in the model, i.e. which groups of animals are to get in contact with each other. For each herd type, a contact structure is defined once for the indoor ( $P_i(ij)$ ) and once for the outdoor ( $P_o(ij)$ ) period.

In addition to the contact structure, herd neighbourhood connections are determined during initialisation. For each herd  $i$ , all other herds  $j$  are assigned to be neighbouring herds for which distance  $d_{ij} \leq r_{neigh}$ . These herds are then added to the state variable neighbours in the *HerdBohvNeighbors* component.

### 1.3.4 Submodels

#### 1.3.4.1 Replacement

**Replacement estimation** In the model, each farmer of a breeding herd estimates how many of its pregnant heifers he is going to keep for replacement of older cows. At the beginning of the simulation, each herd is assigned a random time offset for the estimation, taking into account the time interval ( $\tau_{est}$ ) between the replacement estimates:

$$o_{est} = \lfloor U([0, \tau_{est})) \rfloor \quad (1.1)$$

At every model step ( $t$ ), replacement estimation is performed for all herds where

$$t \bmod \tau_{est} == o_{est} \quad (1.2)$$

Then, for each herd to estimate in a step, the probability to keep pregnant heifers  $p_{keep}$  is

Table 1.4: Model parameters related to replacement

Parameter	Symbol	Type	Description	Value
estimationInterval	$\tau_{est}$	int	Time interval between replacement estimates (weeks)	XX
ticksBeforeCalving	$\tau_{beforeCalving}$	int	How many weeks before calving to make replacement decision	XX
sellTicksBeforeCalving	$\tau_{sell}$	int	How many weeks before calving to sell	XX

estimated based on herd type, target breeders  $n_{targ}$  and number of cows  $n_{cow}$  and bred heifers  $n_{hb}$ . The number of required reproducing animals is

$$n_{req} = n_{targ} - n_{cow} \quad (1.3)$$

This number is used then to calculate a farm's probability to keep pregnant heifers as

$$p_{keep} = \frac{n_{req}}{n_{hb}} \quad (1.4)$$

The result is written to the *HerdManagement* component.

**Replacement decision** Based on the replacement estimate, a replacement decision operator decides for individual bred heifers to keep or sell them. For an individual animal the decision is made when

$$t == t_{conc} + \tau_{gest} - \tau_{est} \quad (1.5)$$

where  $t$  is the current model step,  $t_{conc}$  is the date of conception,  $\tau_{gest}$  is the gestation length and  $\tau_{est}$  a parameter determining how many weeks before calving to make the decision.

Each animal to decide for is stochastically kept with the herd's probability  $p_{keep}$ . If an animal is to be sold, a *ScheduledForSale* component is attached to the animal and the intended date of sale is calculated as

$$t_{sell} = t_{conc} + \tau_{gest} - \tau_{sell} \quad (1.6)$$

while  $\tau_{sell}$  is a model parameter determining how many weeks before calving the animal is to be sold.

#### 1.3.4.2 Grouping

#### 1.3.4.3 Breeding

All animals eligible for breeding are subject to a breeding process that determines the future time point of conception. Cows and Heifers eligible for breeding are inseminated when the herd is in its breeding season (see *HerdManagement* component). Cows are bred if breeding success has not yet been determined and they have exceeded the waiting time after their last calving:

$$t \geq t_{conc} + \tau_{gest} + \tau_{wait}^{(?)} \quad (1.7)$$

where  $t$  is the model step,  $t_{conc}$  is the date of last conception,  $\tau_{gest}$  is gestation length and  $\tau_{wait}^{(?)}$  is the herd type dependent waiting time (either  $w_D$  or  $w_B$ ). Heifers are eligible for breeding if their age is at least  $minAgeBreeding$ .

All eligible animals are inseminated and it is decided stochastically by drawing from a predetermined herd dependent probability ( $f_D$  or  $f_B$ ) if breeding will be successful.

If breeding is successful, the state variable `inseminationSuccess` in the *AwaitsBreeding* component is set to *true* and the future time point of conception/successful insemination is drawn from the herd type dependent distribution  $cd_D$  or  $cd_B$ :

$$t_{conc} = t + bin \cdot \tau_{bins} + [U([0, \tau_{bin}))] \quad (1.8)$$

where  $bin$  is an index of the distribution  $cd?$ , drawn with probabilities proportional to the values in the distribution list.

Animals with breeding success are converted to pregnant when the model step equals their insemination tick:  $t == t_{conc}$ . The *AwaitsBreeding* component is detached and the *IsPregnant* component is attached. The state variables `dateOfConception` from the *AwaitsBreeding* component is transferred to the *IsPregnant* component. Sex of the fetus is determined stochastically according  $p_{female}$ . Breed of the fetus is taken from the dam's breed, except for dairy herds.

For Irish dairy herds, artificial insemination has become the breeding norm on the majority of dairy farms in the Republic of Ireland. But also the breeding bull (a beef bull used for breeding purposes) remains an integral part of the breeding process in many farms. It is used after the breeding season to inseminate cows that have not become pregnant through artificial insemination. To cover this in the model, we have extended the breeding season of all dairy farms by the parameter *dairyBullBreedingDuration*. The fetus of all dairy animals that conceive during this extended period will be mixed, otherwise the dairy breed is assigned.

If the inseminated animal is in Cohort Heifer, it's Cohort is changed to HeiferBred. Animals without breeding success are converted to Cohort Fattener. Further, component *AwaitsBreeding* is detached.

#### 1.3.4.4 Abortion

In the model, pregnant animals can abort their fetus. After successful breeding, abortion occurs with probability  $p_{abort}$ . In case of abortion, the time point is drawn uniformly distributed over the gestation length:

$$t_{abort} = t_{conc} + U([0, \tau_{gest}]) \quad (1.9)$$

where  $t_{conc}$  is the date of conception and  $\tau_{gest}$  is the gestation length. When the date of abortion

$t_{abort}$  equals the current model tick, the *IsPregnant* component is removed from the entity and the *AwaitsBreeding* component is attached. Pre-calculation of abortion date is done for performance. The alternative way of stochastic evaluation of abortion per model step would require the generation of large amount of random numbers, which is computationally expensive.

Table 1.5: Model parameters related to abortion

Parameter	Symbol	Type	Description	Value
abortionProb	$p_{abort}$	float	Probability that a pregnant animal will abort during pregnancy	XX

### 1.3.4.5 Calving

In the model, each pregnant animal gives birth to a calf when model step

$$t = t_{conc} + \tau_{gest} \quad (1.10)$$

where  $t_{conc}$  is the date of conception of the pregnant animal and  $\tau_{gest}$  is the gestation length.

Sex and breed of the calf are taken from *IsPregnant.sex* and *IsPregnant.breed* (set at successful breeding by the breeding submodel). The *AnimalBase* state variable *dateOfBirth* is set to the current model tick. *AnimalGroup.herd* is set to the dam's herd, and the calf is added to the herds list of animals (*HerdBase.animals*).

The BoHV-1 status of the calf depends on the BoHV-1 status of the dam. Generally, each newborn animal is born susceptible, but it can have maternally derived antibodies (*AnimalBohvEpiStat.hasMab == true*) if the dam is either primary or secondary infected during birth. Latently infected animals can also transmit maternally derived antibodies if their last epidemiological status change is smaller than  $\tau_{durationMAB}$ . All animals with maternally derived antibodies are seropositive.

Table 1.6: Model parameters related to calving

Parameter	Symbol	Type	Description	Value
durationMAB	$\tau_{durationMAB}$	float	Duration during which an animal transmits MAB following an epi. status change	XX

### 1.3.4.6 Transports

Animal transports emerge from supply and demand, which emerges from the herd type specific management practices. At the animal level, the process of trade consists of multiple stages. Whenever an animal is intended to leave a herd through the herd type specific management processes, a *ScheduledForSale* component is added to the animal giving information on the intended date of sale. When the date is approached the animal gets the *IsForSale* component making the animal enter a virtual open market. If the animal is not sold for a certain time, the animal is removed. The decision to which herd an animal is moved is managed by two submodels (demand & supply) that are explained in the following.

**Demand** At the beginning of the simulation, each herd is assigned a random week of the year (1-52) at which the herd is to estimate its demand. Based on this initial time, the herd will continuously calculate its demand with the frequency of the estimation interval.

For each herd to estimate in a step, the absolute demand for trade cohorts is estimated based on herd type, target size/breeders and the total number of animals or cows and bred heifers. For breeding herds the demand ( $D_{breeding}$ ) is estimated as

$$D_{breeding} = n_{target} - (n_{cows} + n_{bredHeifers}) \quad (1.11)$$

while  $n_{target}$  is the desired number of target breeders (given in the initialization table) and  $n_{cows}$  and  $n_{bredHeifers}$  the current number of cows and bred heifers, respectively.

Demand of non-breeding herds is calculated as

$$D_{nonBreeding} = n_{target} - n_{totalAnimals} \quad (1.12)$$

while  $totalAnimals$  refers to the total number of animals present in the herd while demand estimation.

A tolerance parameter is accounted for, targeting to produce fewer movements, but larger batches of animals, resulting in fewer trade partners per herd.

**Supply** In the model, supply of animals is generated by the herd type specific management rules (see Table????). Animals are marked there as scheduled for sale and get the component *ScheduledForSale* attached. In this component the future tick of sale is given. When the date is approached the *ScheduledForSale* component is detached and the animal gets the *IsForSale* component. Then, the trading cohort is assigned based on the herd type of the source herd and an animal's sex and pregnancy status.

#### 1.3.4.7 Infection

In the model, each individual animal is assigned to one of four BoHV-1 epidemiological states (see Figure 1.3). Healthy animals that have never been exposed to BoHV-1 are susceptible animals. If a susceptible animal becomes infected it enters the primary infection state and will be a lifelong carrier of the virus. The infectious period of a primary infected animal is two weeks. During this period the animal sheds high levels of virus and is highly contagious. After primary infection hosts recover and enter the latent infection health-state. Latently infected animals are still BoHV-1 positive but do not shed the virus. Latently infected animals can reactivate the virus, which causes a secondary infection. In the model the secondary infection period lasts for one week and affected individuals again pose a risk of infection to susceptible animals. After recovery from secondary infection, animals enter the latent infection health-state.




Figure 1.3: BoHV-1 epidemiological states in the model.

In addition to the epidemiological states, animals can have a certain level of protection due to maternal antibodies or vaccination. Even if animals have these level of protection, they can still become infected with the virus. In the model, these animals do also excrete virus, but to a lesser extent. Also the probability to which seropositive animals reactivate the virus depends on the level of protection.

**Virus transmission** The event of getting infected with BoHV-1 is modelled stochastically by means of a frequency-dependent transmission rate that is converted into an individual probability of getting infected. The frequency-dependent transmission rate accounts for the event of contracting the infection via direct contact and via the neighbourhood.

In the model animals of a herd are split into separate pastures. We assume that infection occurs within and between these pastures by primary and secondary infected animals. First, each herd counts its numbers of primary infected and secondary infected animals (including the vaccinated and those with maternal antibodies), total animals and seropositives, per pasture and in total, and writes them to the state variables of its `HerdBohvStatus` component. These are in the next stage used to calculate infection probabilities.

Then, each herd stochastically determines individuals to be infected according the following scheme: The herd sums up the number of PIs and total animal number in it's neighboring herds  $k$  for neighborhood infection.

Next, the herd iterates over all possible Pasture-to-Pasture combinations  $ij$  and calculates the individual probability of infection for each pasture  $i$  from animals in pasture  $j$  as a combination of three frequency-dependent transmission rates:

$$P_{inf,i} = 1 - \exp^{-(withinPool+betweenPool+betweenHerds)} \quad (1.13)$$

The *withinPool* transmission rate describes the rate at which an individual receives the infection from its own pasture as follows, while  $pi_j$ ,  $si_j$ ,  $piProtect_j$ ,  $siProtect_j$  and  $n_j$  are the number of

primary (protect) infected, secondary (protect) infected and total animals in pasture  $j$ .

$$\begin{aligned}
 \text{withinPool} = & \beta_{pi}^{(i)} \frac{\sum_j^{j=i \wedge p_{ij}} p_{ij}}{\sum_j^{j=i \wedge p_{ij}} n_j} + \beta_{si}^{(i)} \frac{\sum_j^{j=i \wedge p_{ij}} s_{ij}}{\sum_j^{j=i \wedge p_{ij}} n_j} + \\
 & \beta_{piProtect}^{(i)} \frac{\sum_j^{j=i \wedge p_{ij}} piProtect_j}{\sum_j^{j=i \wedge p_{ij}} n_j} + \\
 & \beta_{siProtect}^{(i)} \frac{\sum_j^{j=i \wedge p_{ij}} siProtect_j}{\sum_j^{j=i \wedge p_{ij}} n_j}
 \end{aligned} \tag{1.14}$$

Transmission between different management groups (between cohort transmission) is only triggered by unprotected primary and secondary infected animals, calculated as follows:

$$\text{betweenPool} = \beta_{pi}^{(p)} \cdot \frac{\sum_j^{j \neq i \wedge \neg p_{ij}} p_{ij}}{\sum_j^{j \neq i \wedge \neg p_{ij}} n_j} + \beta_{si}^{(p)} \cdot \frac{\sum_j^{j \neq i \wedge \neg p_{ij}} s_{ij}}{\sum_j^{j \neq i \wedge \neg p_{ij}} n_j} \tag{1.15}$$

In addition, between herd infection is modelled as:

$$\text{betweenHerds} = \beta_{pi}^{(h)} \frac{p_{i\_neigh}^i}{n_{neigh}} \tag{1.16}$$

Here,  $p_{i\_neigh}^i$  describes all primary infected animals in the neighbouring herds and  $n_{neigh}$  all total animals in the neighbouring herds.

In a last step, the herd iterates over all susceptible animals in all pastures  $i$  where  $P_{inf,i} > 0$ , and infects each individual with probability  $P_{inf,i}$ . Infected individuals are converted to primary infected animals.

**Virus reactivation** Latently infected animals have the potential to reactivate the virus which causes a secondary infection. Virus reactivation is simulated as a binomial random process, with a constant weekly reactivation probability calculated as

$$P_{reactivate} = 1 - (1 - r_{reactivattion})^{1/52} \tag{1.17}$$

with  $r_{reactivattion}$  being the annual reactivation rate for animals in the latent infection state. The reactivation rate differs, depending on the protection level of the animals. Vaccinated animals or those with maternal antibodies have a lower risk of reactivating the virus, compared to those with no level of protection. Animals transports increases the reactivation probability by orders of magnitude.

**Recovery** Primary infected animals convert to latently infected animals  $\tau_{primaryInfected}$  steps after infection. Secondary infected animals remain  $\tau_{secondaryInfected}$  steps infectious before they become latently infected again. Animals with maternal antibodies loose their level of protection, if not vaccinated, when their age equals  $\tau_{secondaryInfected} + 1$  (in weeks).

### 1.3.4.8 Mortality

The model accounts for a background mortality, representing the constant weekly risk that an individual animal will die a natural death. Instead of evaluating mortality at each simulation step, life expectancy is calculated for each animal (whenever an animal is added to the system/model) based on the annual background mortality parameter  $r_{mortality}$  as

$$LifeExpectancy = - \left( \frac{\log(x)}{r_{mortality}} \right) \quad (1.18)$$

with  $x$  being a random number between 0 and 1. If the calculated *LifeExpectancy* exceeds a predefined threshold (see Table 1.1) a new life expectancy is calculated by a means of a new  $r_{mortality}$ . The  $r_{mortality}$  parameter is chosen depending on the belonging of an animal to a certain age class. The date of death is calculated accordingly and set in the *AnimalBase* component. If the current model step equals an animals date of death, the animal is removed from the engine.

Table 1.7: Model parameters related to mortality

Parameter	Symbol	Type	Description	Value
backgroundMortalityNewborn	$r_{newborn}$	float	Annual mortality rate of newborn animals	0.87
backgroundMortalityCalf	$r_{calf}$	float	Annual mortality rate of calves	0.05
backgroundMortalityYoungstock	$r_{youngstock}$	float	Annual mortality rate of youngstock	0.05
backgroundMortalityAdults	$r_{adults}$	float	Annual mortality rate of adults	0.01
backgroundMortalityOldAdultsDairy	$r_{oldAdultsDairy}$	float	Annual mortality rate of old dairy adults	0.3
backgroundMortalityOldAdultsBeef	$r_{oldAdultsBeef}$	float	Annual mortality rate of old beef adults	0.05
backgroundMortalityFattener	$r_{fattener}$	float	Annual mortality rate of fatteners	3

### 1.3.4.9 Virus release

In the model, we released the virus by setting animals over 3 years old to latently infected in 60% of all herds (randomly selected). This was done to ensure a valid age-dependent prevalence in the herds and to reduce the burn-in period after virus release.

# Bibliography

---

- Brock, J., Lange, M., More, S., Graham, D., & Thulke, H. (2020). Reviewing age-structured epidemiological models of cattle diseases tailored to support management decisions: Guidance for the future. *Preventive Veterinary Medicine*, 174(October 2019), 104814.
- DAFM. (2017). *AIM Statistics Report 2017 (2018)* (Tech. Rep.). Retrieved from <https://www.agriculture.gov.ie/media/migration/animalhealthwelfare/animalidentificationandmovement/AIMStatisticsReport2017260618.pdf>
- Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V., Giske, J., ... DeAngelis, D. (2006). A standard protocol for describing individual-based and agent-based models. *Ecological Modelling*, 198(1-2), 115–126.
- Grimm, V., Berger, U., DeAngelis, D., Polhill, J., Giske, J., & Railsback, S. (2010). The ODD protocol: A review and first update. *Ecological Modelling*, 221(23), 2760–2768.
- Grimm, V., Railsback, S., Vincenot, C., Berger, U., Gallagher, C., DeAngelis, D., ... Ayllón, D. (2020). The ODD Protocol for Describing Agent-Based and Other Simulation Models: A Second Update to Improve Clarity, Replication, and Structural Realism. *Journal of Artificial Societies and Social Simulation*, 23(2), 7.